


SIDDHARTH GROUP OF INSTITUTIONS :: PUTTUR

Siddharth Nagar, Narayanavanam Road – 517583

QUESTION BANK (DESCRIPTIVE)
Subject with Code: MATHEMATICS-III(18HS0834)

Branch: B.Tech(ECE)

Year & Sem: II-B.Tech&I-Sem

UNIT -I
Regulation: R18

NUMERICAL METHOD -I
1.a) Write the formula to find the root of an equation by Regula Falsi method [2M]

b) Write Simpson formulae [2M]

c) Write the formula to find a cube root of a number by Newton Raphson's method [2M]

d) Evaluate $\Delta \tan^{-1} x$ [2M]

e) Construct a forward difference table for the function $y = x^3$ for $x = 0, 1, 2, 3, 4, 5$. [2M]

2. Find a positive root of $x^3 - x - 1 = 0$ correct to two decimal places by Bisection method. [10 M]

3. Find out the root of the equation $x \log_{10}(x) = 1.2$ using False position method. [10 M]

4. Find the root of the equation $xe^x = 2$ using Regula-falsi method. [10 M]

5. Find a real root of the equation $xe^x - \cos x = 0$ using Newton-Raphson method. [10 M]

6. Using Newton-Raphson Method

(i) Find square root of 28.

(ii) Find cube root of 15.

[10M]

7. From the following table values of x and

 $y = \tan x$ interpolate values of y when

 $x = 0.12$ and $x = 0.28$

x	0.10	0.15	0.20	0.25	0.30
y	0.1003	0.1511	0.2027	0.2553	0.3093

[10M]

8.a) Using Newton's forward interpolation formula and the given table of values

x	1.1	1.3	1.5	1.7	1.9
$f(x)$	0.21	0.69	1.25	1.89	2.61

 Obtain the value of $f(x)$ when $x = 1.4$

[5M]

b) Use Newton's Backward interpolation formula to find $f(32)$

 given $f(25) = 0.2707$, $f(30) = 0.3027$, $f(35) = 0.3386$, $f(40) = 0.3794$

[5M]

9. Evaluate $\int_0^1 \frac{1}{1+x} dx$ **(i)** By trapezoidal rule and Simpson's $\frac{1}{3}$ rule

[5M]

(ii) Using Simpson's $\frac{3}{8}$ rule and compare the result with actual value

[5M]

10.a) Compute $\int_0^4 e^x dx$ by Simpson's $\frac{1}{3}$ rule with 10 sub divisions.

[5M]

b) Compute $\int_3^7 x^2 \log x dx$ using trapezoidal rule and Simpson's rule by taking 10 sub divisions. [5M]

[Type text]



SIDDHARTH GROUP OF INSTITUTIONS :: PUTTUR
Siddharth Nagar, Narayanavanam Road – 517583

QUESTION BANK (DESCRIPTIVE)

Subject with Code :MATHEMATICS-III(18HS0834)
Year &Sem: II-B.Tech& I-Sem

Branch: B.Tech(ECE)
Regulation: R18

UNIT –II

NUMERICAL METHOD-II

1. a) write R-K method of 4th order formula [2M]
b) write the diagonal five-point formula [2M]
c) write the Taylor's series solution of $y' = -xy, y(0)=1$ upto x^4 [2M]
d) write the standard five-point formula [2M]
e) Use Euler's method to find $y(0.1)$ given $y' = (x^3 + xy^2)e^{-x}, y(0) = 1$ [2M]
2. a) Tabulate $y(0.1), y(0.2),$ and $y(0.3)$ using Taylor's series method given that $y' = y^2 + x$ and $y(0) = 1$ [5M]
b) Using Euler's method, find an approximate value of y corresponding to $x = 1$ given that $\frac{dy}{dx} = x + y$ and $y = 1$ when $x = 0$. [5M]
- 3 Using Taylor's series method find an approximate value of y at $x = 0.2$ for the D.E $y' - 2y = 3e^x, y(0) = 0$. Compare the numerical solution obtained with exact solution. [10M]
- 4a) Solve $y' = x + y$, given $y(1)=0$ find $y(1.1)$ and $y(1.2)$ by Taylor's series method [5M]
b) Solve by Euler's method [5M]
 $\frac{dy}{dx} = \frac{2y}{x}$ given $y(1) = 2$ and find $y(2)$.
- 5 Using R-K method of 4th order, solve $\frac{dy}{dx} = \frac{y^2 - x^2}{y^2 + x^2}, y(0)=1$ Find $y(0.2)$ and $y(0.4)$ [10M]
- 5 Using R-K method of 4th order find $y(0.1), y(0.2)$ and $y(0.3)$ given that $\frac{dy}{dx} = 1 + xy, y(0) = 2$ [10M]
- 7) a) Using Runge-Kutta method of fourth order, compute $y(0.2)$ from $y' = xy$ $y(0)=1$, taking $h=0.2$ [5M]
b) using Euler's method $y' = y^2 + x, y(0)=1$. Find $y(0.1)$ and $y(0.2)$ [5M]


MATHEMATICS-III

[Type text]


8) Solve $y'' - x(y')^2 + y^2 = 0$ using R-K method of 4th order for $x=0.2$ given $y(0) = 1, y'(0)=0$ taking $h=0.2$
[10M]

9)) Solve the Laplace Equation $u_{xx} + u_{yy} = 0$ given that,

[10M]



10) Evaluate the function $u(x, y)$ satisfying $\nabla^2 u = 0$ at the pivotal points given the boundary values as follows:



[10M]

[Type text]



SIDDHARTH GROUP OF INSTITUTIONS :: PUTTUR

Siddharth Nagar, Narayanavanam Road – 517583

QUESTION BANK (DESCRIPTIVE)

Subject with Code : MATHEMATICS-III(18HS0834)

Branch: B.Tech(ECE)

Year & Sem: II-B.Tech& I-Sem

Regulation: R18

TRANSFORMS CALCULUS-I

UNIT-III

1. a) Find the Laplace transform of $e^{at} \cosh bt$ [2 M]
b) Find the Laplace transform of $3 \cos 3t \cdot \cos 4t$ [2M]
- c) Find $L^{-1} \left\{ \frac{2s-5}{4s^2+25} \right\}$ by using linear property. [2 M]
- d) Find $L \{t^2 + 3t + 10\}$ [2M]
e) State Convolution theorem. [2M]
2. a) Find the Laplace transform of $e^{-3t} (2 \cos 5t - 3 \sin 5t)$ [5M]
- b) Find the Laplace transform of $f(t) = \int_0^t e^{-t} \cos t dt$. [5M]
3. a) Find the Laplace transform of $f(t) = \frac{1 - \cos at}{t}$ [5 M]
- b) Show that $\int_0^{\infty} t^2 e^{-4t} \cdot \sin 2t dt = \frac{11}{500}$, Using Laplace transform [5 M]
4. a) Find the Laplace Transform of Square-wave function of period $2a$,
defined as $f(t) = \begin{cases} k, & 0 < t < a \\ -k, & a < t < 2a \end{cases}$ [5 M]
- b) Using Laplace transform, evaluate $\int_0^{\infty} \frac{\cos at - \cos bt}{t} dt$. [5 M]
5. a) Find the Laplace transform of $f(t) = e^{-4t} \int_0^t \frac{\sin 3t}{t} dt$. [5 M]
- b) Find the Laplace transform of $f(t) = t e^{2t} \sin 3t$ [5 M]
6. a) Find $L^{-1} \left\{ \frac{3s-2}{s^2-4s+20} \right\}$ by using first shifting theorem. [5M]
- b) Find $L^{-1} \left\{ \frac{1}{2} \log \left(\frac{s^2+a^2}{s^2+b^2} \right) \right\}$ [5M]
7. a) Find $L^{-1} \left\{ \frac{1}{(s^2+5^2)^2} \right\}$, using Convolution theorem. [5 M]
- b) Find $L^{-1} \left\{ \frac{s^2}{(s^2+4)(s^2+25)} \right\}$, using Convolution theorem. [5M]

[Type text]

8. a) Find the Inverse Laplace transform of $\frac{1}{s^2(s^2 + a^2)}$. [5 M]

b) Find $L^{-1}\left\{s \log\left(\frac{s-1}{s+1}\right)\right\}$ [5 M]

9. Using Laplace transform method to solve $y^{11} - 3y^1 + 2y = 4t + e^{3t}$ where $y(0) = 1, y^1(0) = 1$ [10M]

10. Solve the D.E $\frac{d^2x}{dt^2} + 9x = \sin t$ using Laplace Transform given that

$x(0) = 1, x\left(\frac{\pi}{2}\right) = 1$ [10M]

UNIT – IV

TRANSFORMS CALCULUS-II

[Type text]

1. a) Define fourier sine and cosine transforms [2M]
b) Find the fourier sine transform of $\frac{1}{x}$ [2M]
c) Define the inverse fourier sine and cosine transforms [2M]
d) Find the fourier cosine transform of e^{-ax} , $a > 0$ and hence deduce the inverse formula [2M]
e) Find the finite fourier sine transform of $f(x) = 2x$, $0 < x < 4$. [2M]

2. a) Express $f(x) = \begin{cases} 1, 0 \leq x \leq \pi \\ 0, x > \pi \end{cases}$ as a fourier sine integral and hence evaluate

$$\int_0^{\infty} \frac{1 - \cos(\pi\lambda)}{\lambda} \sin(x\lambda) d\lambda \quad [5M]$$

- b) Prove that (i) $F_s \{ a f(x) + b g(x) \} = a F_s(p) + b G_s(p)$

(ii) $F_c \{ a f(x) + b g(x) \} = a F_c(p) + b G_c(p)$ [5M]

3. a) Prove that $F[x^n f(x)] = (-i)^n \frac{d^n}{dp^n} [F(p)]$ [5M]

- b) Prove that $F_s \{ x f(x) \} = -\frac{d}{dp} [F_c(p)]$ [5M]

4. Find the Fourier transform of $f(x) = \begin{cases} a^2 - x^2, |x| < a \\ 0, |x| > a > 0 \end{cases}$ Hence show that

$$\int_0^{\infty} \frac{\sin x - x \cos x}{x^3} dx = \frac{\pi}{4} \quad [10M]$$

5. a) Find the Fourier transform of $f(x) = e^{-\frac{x^2}{2}}$, $-\infty < x < \infty$ [5M]

- b) If $F(p)$ is the complex Fourier transform of $f(x)$, then prove that the complex Fourier transform of $f(x) \cos ax$ is $\frac{1}{2} [F(p+a) + F(p-a)]$ [5M]

6. a) Find the Fourier cosine transform of $e^{-ax} \cos ax$, $a > 0$ [5M]

- b) Find the Fourier cosine transform of $f(x) = \begin{cases} x, \text{ for } 0 < x < 1 \\ 2 - x, \text{ for } 1 < x < 2 \\ 0, \text{ for } x > 2 \end{cases}$ [5M]

7. Find the Fourier sine and cosine transforms of $f(x) = \frac{e^{-ax}}{x}$ and deduce that

$$\int_0^{\infty} \frac{e^{-ax} - e^{-bx}}{x} \sin sx dx = \tan^{-1} \left(\frac{s}{a} \right) - \tan^{-1} \left(\frac{s}{b} \right) \quad [10M]$$

8. Find the Fourier sine and cosine transforms of $f(x) = e^{-ax}$, $a > 0$ and hence deduce the integrals

(i) $\int_0^{\infty} \frac{p \sin px}{a^2 + p^2} dp$ (ii) $\int_0^{\infty} \frac{\cos px}{a^2 + p^2} dp$ [10M]

9. Find the inverse Fourier sine transform of $f(x)$ of $F_s(p) = \frac{p}{1 + p^2}$ [10M]

[Type text]

10.a) Find the finite Fourier sine transform of $f(x)$, defined by $\left\{ \begin{array}{l} x, 0 \leq x \leq \frac{\pi}{2} \\ \pi - x, \frac{\pi}{2} \leq x \leq \pi \end{array} \right\}$ [5M]

b) Find the inverse finite Fourier sine transform of $f(x)$, if $F_s(n) = \frac{16(-1)^{n-1}}{n^3}$, where n is a positive integer and $0 < x < 8$.

[5M]

[Type text]



SIDDHARTH GROUP OF INSTITUTIONS :: PUTTUR

Siddharth Nagar, Narayanavanam Road – 517583

QUESTION BANK (DESCRIPTIVE)

Subject with Code : MATHEMATICS-III(18HS0834)

Branch: B.Tech(ECE)

Year & Sem: II-R Tech& I-Sem

Regulation: R18

UNIT – V

PARTIAL DIFFERENTIAL EQUATIONS

1. a) Solve $xp + yq = 3z$. [2 M]
b) Solve $r + 6s + 9t = 0$. [2 M]
c) Solve $p(1 + q) = qz$. [2 M]
d) Solve $\frac{\partial^3 z}{\partial x^3} - 4\frac{\partial^3 z}{\partial x^2 \partial y} + 4\frac{\partial^3 z}{\partial x \partial y^2} = 0$. [2 M]
e) Find the particular integral of the equation $4r + 12s + 9t = e^{3x-2y}$. [2 M]
2. a) Solve $(x^2 - y^2 - z^2)p + 2xyq = 2xz$. [5 M]
b) Solve $(z - y)p + (x - z)q = y - x$. [5 M]
3. a) Solve $x(y - z)p + y(z - x)q = z(x - y)$. [5 M]
b) Solve $x^2(y - z)p + y^2(z - x)q = z^2(x - y)$. [5 M]
4. a) Solve $p^2 + q^2 = x + y$. [5 M]
b) Solve $z^2(p^2 x^2 + q^2) = 1$. [5 M]
5. a) Solve $r - 4s + 4t = e^{2x+y}$. [5 M]
b) Solve $\frac{\partial^2 z}{\partial x^2} + \frac{\partial^2 z}{\partial x \partial y} - 6\frac{\partial^2 z}{\partial y^2} = \cos(2x + y)$. [5 M]
6. a) Solve $(D^2 + 3DD' + 2D'^2)z = 24xy$. [5 M]
b) Solve $\frac{\partial^3 z}{\partial x^3} - 2\frac{\partial^3 z}{\partial x^2 \partial y} = 2e^{2x} + 3x^2y$. [5 M]
7. a) Solve $(D^2 + 2DD' + D'^2 - 2D - 2D')z = \sin(x + 2y)$. [5 M]
b) Solve $(D - D' - 1)(D - D' - 2)z = e^{2x-y}$. [5 M]
8. A tightly stretched string of length l with fixed ends is initially in equilibrium position. It is set vibrating by giving each point a velocity $b \sin^3\left(\frac{\pi x}{l}\right)$. Find the displacement $y(x, t)$. [10 M]
9. A tightly stretched string with fixed end points $x=0$ and $x=l$ is initially at rest in its equilibrium position. It is set vibrating by giving each point a velocity $kx(l - x)$. Find the displacement of the string at any distance x from one end at any time t . [10 M]
10. A homogeneous rod of conducting material of length 100cm has its ends kept at zero temperature and the temperature initially is $u(x, 0) = x, \quad 0 \leq x \leq 50$
 $= 100 - x, \quad 50 \leq x \leq 100$
Find the temperature $u(x, t)$ at any time. [10 M]