



**SIDDHARTHA INSTITUTE OF SCIENCE AND TECHNOLOGY:: PUTTUR
(AUTONOMOUS)**

Siddharth Nagar, Narayanavanam Road – 517583

QUESTION BANK (DESCRIPTIVE)

Subject with Code: NMPS(19HS0833)

Course & Branch: B.Tech - CE

Year & Sem: II-B.Tech &II-Sem

Regulation: R19

**UNIT -I
SOLUTIONS OF ALGEBRAIC AND TRANSCENDENTAL
EQUATIONS, INTERPOLATION**

1	Find out the square root of 25 given $x_0 = 2.0$, $x_1 = 7.0$ using Bisection method.	[L1][CO1]	[12M]												
2	Find a positive root of $x^3 - x - 1 = 0$ correct to two decimal places by Bisection method	[L1][CO1]	[12M]												
3	Find a positive root of $f(x) = e^x - 3$ correct to two decimal places by Bisection method	[L1][CO1]	[12M]												
4	Find a real root of the equation $xe^x - \cos x = 0$ using Newton – Raphson method	[L1][CO1]	[12M]												
5	Using Newton-Raphson method (i) Find square root of 28 (ii) Find cube root of 15	[L3][CO1]	[12M]												
6	a. Using Newton-Raphson method Find reciprocal of 12. b. Find a real root of the equation $x \tan x + 1 = 0$ using Newton – Raphson method	[L3][CO1] [L1][CO1]	[06M] [06M]												
7	Find out the root of the equation $x \log_{10}(x) = 1.2$ using False position method.	[L1][CO1]	[12M]												
8	Find the root of the equation $x e^x = 2$ using Regula-falsi method.	[L1][CO1]	[12M]												
9	From the following table values of x and $y = \tan x$. Interpolate values of y when $x = 0.12$ and $x = 0.28$.	[L1][CO1]	[12M]												
	<table border="1"> <tbody> <tr> <td>x</td> <td>0.10</td> <td>0.15</td> <td>0.20</td> <td>0.25</td> <td>0.30</td> </tr> <tr> <td>y</td> <td>0.1003</td> <td>0.1511</td> <td>0.2027</td> <td>0.2553</td> <td>0.3093</td> </tr> </tbody> </table>	x	0.10	0.15	0.20	0.25	0.30	y	0.1003	0.1511	0.2027	0.2553	0.3093		
x	0.10	0.15	0.20	0.25	0.30										
y	0.1003	0.1511	0.2027	0.2553	0.3093										
10	a. Using Newton's forward interpolation formula and the given table of value Obtain the value of $f(x)$ when $x = 1.4$.	[L3][CO1]	[06M]												
	<table border="1"> <tbody> <tr> <td>x</td> <td>1.1</td> <td>1.3</td> <td>1.5</td> <td>1.7</td> <td>1.9</td> </tr> <tr> <td>$f(x)$</td> <td>0.21</td> <td>0.69</td> <td>1.25</td> <td>1.89</td> <td>2.61</td> </tr> </tbody> </table>	x	1.1	1.3	1.5	1.7	1.9	$f(x)$	0.21	0.69	1.25	1.89	2.61		
x	1.1	1.3	1.5	1.7	1.9										
$f(x)$	0.21	0.69	1.25	1.89	2.61										
	b. Use Newton's backward interpolation formula to find $f(32)$ given $f(25) = 0.2707$, $f(30) = 0.3027$, $f(35) = 0.3386$, $f(40) = 0.3794$.	[L3][CO1]	[06M]												

UNIT –II
NUMERICAL SOLUTIONS OF ORDINARY DIFFERENTIAL
EQUATIONS, NUMERICAL INTEGRATIONS

1	Tabulate $y(0.1)$, $y(0.2)$ and $y(0.3)$ using Taylor's series method given that $y' = y^2 + x$ and $y(0) = 1$	[L6][CO2]	[12M]
2	Using Taylor's series method find an approximate value of y at $x = 0.2$ for the D.E $y' - 2y = 3e^x$, $y(0) = 0$. Compare the numerical solution obtained with exact solution.	[L3][CO2]	[12M]
3	a. Solve $y' = x + y$, given $y(1) = 0$ find $y(1.1)$ and $y(1.2)$ by Taylor's series method. b. Solve by Euler's method $\frac{dy}{dx} = \frac{2y}{x}$ given $y(1) = 2$ and find $y(2)$.	[L3][CO2] [L3][CO2]	[06M] [06M]
4	Using Euler's method, find an approximate value of y corresponding to $x = 1$ given that $\frac{dy}{dx} = x + y$ and $y = 1$ when $x = 0$ taking step size $h = 0.1$.	[L3][CO2]	[12M]
5	a. Using Euler's method $y' = y^2 + x$, $y(0) = 1$. Find $y(0.1)$ and $y(0.2)$ b. Using Runge – Kutta method of fourth order, compute $y(0.2)$ from $y' = xy$, $y(0) = 1$, taking $h = 0.2$	[L3][CO2] [L3][CO2]	[06M] [06M]
6	Using R-K method, evaluate $y(0.1)$ and $y(0.2)$ given $y' = x + y$; $y(0) = 1$.	[L3][CO2]	[12M]
7	Using R-K method of 4 th order find $y(0.1)$, $y(0.2)$ and $y(0.3)$ given that $\frac{dy}{dx} = 1 + xy$, $y(0) = 2$.	[L3][CO2]	[12M]
8	Solve $y'' - x(y')^2 + y^2 = 0$ using R-K method of 4 th order for $x = 0.2$ given $y(0) = 1$, And $y'(0) = 0$ taking $h = 0.2$	[L3][CO2]	[12M]
9	Evaluate $\int_0^1 \frac{1}{1+x} dx$ (i) by Trapezoidal rule and Simpson's $\frac{1}{3}$ -rule. (ii) using Simpson's $\frac{3}{8}$ rule and compare the result with actual value.	[L5][CO2]	[12M]
10	a. Compute $\int_0^4 e^x dx$ by Simpson's $\frac{3}{8}$ rule with 12 sub divisions. b. Compute $\int_3^7 x^2 \log x dx$ by Trapezoidal rule and Simpson's $\frac{1}{3}$ rule by taking 10 sub divisions.	[L5][CO2] [L5][CO2]	[06M] [06M]

UNIT-III
BASIC STATISTICS&BASIC PROBABILITY

1	a)i) The weights of 6 competitors in a game are given below 58,62,56,63,55,61 kgs. Find arithmetic mean of weight of competitors. ii) Find the median of the following values 26, 8, 6, 12, 15, 32.	[L3][CO3]	[06M]																																		
	b. Find arithmetic mean to the following data using step deviation method <table border="1" style="margin: 10px auto; border-collapse: collapse;"> <tr> <td style="padding: 2px;">Marks</td> <td style="padding: 2px;">10-20</td> <td style="padding: 2px;">20-30</td> <td style="padding: 2px;">30-40</td> <td style="padding: 2px;">40-50</td> <td style="padding: 2px;">50-60</td> </tr> <tr> <td style="padding: 2px;">Frequency</td> <td style="padding: 2px;">5</td> <td style="padding: 2px;">8</td> <td style="padding: 2px;">25</td> <td style="padding: 2px;">22</td> <td style="padding: 2px;">10</td> </tr> </table>	Marks	10-20	20-30	30-40	40-50	50-60	Frequency	5	8	25	22	10	[L1][CO3]	[06M]																						
Marks	10-20	20-30	30-40	40-50	50-60																																
Frequency	5	8	25	22	10																																
2	a. Find the median to the following data <table border="1" style="margin: 10px auto; border-collapse: collapse;"> <tr> <td style="padding: 2px;">Class intervals</td> <td style="padding: 2px;">40-50</td> <td style="padding: 2px;">50-60</td> <td style="padding: 2px;">60-70</td> <td style="padding: 2px;">70-80</td> <td style="padding: 2px;">80-90</td> </tr> <tr> <td style="padding: 2px;">Frequency</td> <td style="padding: 2px;">5</td> <td style="padding: 2px;">12</td> <td style="padding: 2px;">23</td> <td style="padding: 2px;">8</td> <td style="padding: 2px;">2</td> </tr> </table> b. Find arithmetic mean to the following data <table border="1" style="margin: 10px auto; border-collapse: collapse;"> <tr> <td style="padding: 2px;">x</td> <td style="padding: 2px;">1</td> <td style="padding: 2px;">2</td> <td style="padding: 2px;">3</td> <td style="padding: 2px;">4</td> <td style="padding: 2px;">5</td> </tr> <tr> <td style="padding: 2px;">f</td> <td style="padding: 2px;">5</td> <td style="padding: 2px;">8</td> <td style="padding: 2px;">10</td> <td style="padding: 2px;">12</td> <td style="padding: 2px;">6</td> </tr> </table>	Class intervals	40-50	50-60	60-70	70-80	80-90	Frequency	5	12	23	8	2	x	1	2	3	4	5	f	5	8	10	12	6	[L1][CO3]	[06M]										
Class intervals	40-50	50-60	60-70	70-80	80-90																																
Frequency	5	12	23	8	2																																
x	1	2	3	4	5																																
f	5	8	10	12	6																																
3	a) Find mode to the following data <table border="1" style="margin: 10px auto; border-collapse: collapse;"> <tr> <td style="padding: 2px;">X</td> <td style="padding: 2px;">0-5</td> <td style="padding: 2px;">5-10</td> <td style="padding: 2px;">10-15</td> <td style="padding: 2px;">15-20</td> <td style="padding: 2px;">20-25</td> <td style="padding: 2px;">25-30</td> <td style="padding: 2px;">30-35</td> <td style="padding: 2px;">35-40</td> </tr> <tr> <td style="padding: 2px;">F</td> <td style="padding: 2px;">5</td> <td style="padding: 2px;">7</td> <td style="padding: 2px;">10</td> <td style="padding: 2px;">18</td> <td style="padding: 2px;">20</td> <td style="padding: 2px;">12</td> <td style="padding: 2px;">8</td> <td style="padding: 2px;">2</td> </tr> </table> b) Find the median to the following data <table border="1" style="margin: 10px auto; border-collapse: collapse;"> <tr> <td style="padding: 2px;">X</td> <td style="padding: 2px;">5</td> <td style="padding: 2px;">8</td> <td style="padding: 2px;">11</td> <td style="padding: 2px;">14</td> <td style="padding: 2px;">17</td> <td style="padding: 2px;">20</td> <td style="padding: 2px;">23</td> </tr> <tr> <td style="padding: 2px;">F</td> <td style="padding: 2px;">2</td> <td style="padding: 2px;">8</td> <td style="padding: 2px;">12</td> <td style="padding: 2px;">20</td> <td style="padding: 2px;">10</td> <td style="padding: 2px;">6</td> <td style="padding: 2px;">3</td> </tr> </table>	X	0-5	5-10	10-15	15-20	20-25	25-30	30-35	35-40	F	5	7	10	18	20	12	8	2	X	5	8	11	14	17	20	23	F	2	8	12	20	10	6	3	[L1][CO3]	[06M]
X	0-5	5-10	10-15	15-20	20-25	25-30	30-35	35-40																													
F	5	7	10	18	20	12	8	2																													
X	5	8	11	14	17	20	23																														
F	2	8	12	20	10	6	3																														
4	a) Obtain mode of the values 10,12,15,20,12,16,18,15,12,10,16,20,12,24. b) The first four moments of a distribution about the value 5 of the variables are 2, 20, 40 and 50. Calculate mean, variance, β_1 and β_2 of the distribution.	[L1][CO3] [L1][CO3]	[06M] [06M]																																		
5	Compute Karl Pearson and Bowley's coefficient of Skewness to the following data <table border="1" style="margin: 10px auto; border-collapse: collapse;"> <tr> <td style="padding: 2px;">Class intervals</td> <td style="padding: 2px;">0-10</td> <td style="padding: 2px;">10-20</td> <td style="padding: 2px;">20-30</td> <td style="padding: 2px;">30-40</td> <td style="padding: 2px;">40-50</td> <td style="padding: 2px;">50-60</td> <td style="padding: 2px;">60-70</td> <td style="padding: 2px;">70-80</td> <td style="padding: 2px;">80-90</td> <td style="padding: 2px;">90-100</td> </tr> <tr> <td style="padding: 2px;">frequency</td> <td style="padding: 2px;">2</td> <td style="padding: 2px;">6</td> <td style="padding: 2px;">11</td> <td style="padding: 2px;">20</td> <td style="padding: 2px;">40</td> <td style="padding: 2px;">75</td> <td style="padding: 2px;">45</td> <td style="padding: 2px;">25</td> <td style="padding: 2px;">18</td> <td style="padding: 2px;">8</td> </tr> </table>	Class intervals	0-10	10-20	20-30	30-40	40-50	50-60	60-70	70-80	80-90	90-100	frequency	2	6	11	20	40	75	45	25	18	8	[L6][CO3]	[12M]												
Class intervals	0-10	10-20	20-30	30-40	40-50	50-60	60-70	70-80	80-90	90-100																											
frequency	2	6	11	20	40	75	45	25	18	8																											
6	Compute the first four central moments to the following data and also find Sheppard's correction, β_1 and β_2 <table border="1" style="margin: 10px auto; border-collapse: collapse;"> <tr> <td style="padding: 2px;">Class intervals</td> <td style="padding: 2px;">0-10</td> <td style="padding: 2px;">10-20</td> <td style="padding: 2px;">20-30</td> <td style="padding: 2px;">30-40</td> <td style="padding: 2px;">40-50</td> <td style="padding: 2px;">50-60</td> <td style="padding: 2px;">60-70</td> </tr> <tr> <td style="padding: 2px;">Frequency</td> <td style="padding: 2px;">2</td> <td style="padding: 2px;">8</td> <td style="padding: 2px;">12</td> <td style="padding: 2px;">40</td> <td style="padding: 2px;">20</td> <td style="padding: 2px;">15</td> <td style="padding: 2px;">3</td> </tr> </table>	Class intervals	0-10	10-20	20-30	30-40	40-50	50-60	60-70	Frequency	2	8	12	40	20	15	3	[L6][CO3]	[12M]																		
Class intervals	0-10	10-20	20-30	30-40	40-50	50-60	60-70																														
Frequency	2	8	12	40	20	15	3																														
7	a) Three students A,B,C are in running race. A and B have the same Probability of winning and each is twice as likely to win as C. Find the Probability that B or C wins. b) Determine (i) $P\left(\frac{B}{A}\right)$ (ii) $P\left(\frac{A}{B^c}\right)$ if A and B are events with $P(A) = \frac{1}{3}$, $P(B) = \frac{1}{4}$, $P(A \cup B) = \frac{1}{2}$.	[L6][CO3] [L5][CO3]	[06M] [06M]																																		
8	a) In a certain town 40% have brown hair, 25% have brown eyes and 15% have both brown hair and brown eyes. A person is selected at random from the town. i) If he has brown hair, what is the probability that he has brown eyes also? ii) If he has brown eyes, determine the probability, that he does not have brown hair?	[L5][CO3]	[06M]																																		

	b) The probability that students A, B, C, D solve the problem are $\frac{1}{3}$, $\frac{2}{5}$, $\frac{1}{5}$ and $\frac{1}{4}$ respectively. If all of them try to solve the problem, what is the probability that the problem is solved.	[L6][CO3]	[06M]
9	Two dice are thrown. Let A be the event that the sum of the point on the faces is 9. Let B be the event that at least one number is 6. Find (i) $P(A \cap B)$ (ii) $P(A \cup B)$ (iii) $P(A^c \cup B^c)$ (iv) $P(A^c \cap B^c)$ (v) $P(A \cap B^c)$	[L1][CO3]	[12M]
10	In a certain college 25% of boys and 10% of girls are studying mathematics. The girls constitute 60% of the student body. (a) What is the probability that mathematics is being studied? (b) If a student is selected at random and is found to be studying mathematics, find the probability that the student is a girl? (c) a boy.	[L6][CO3]	[12M]

**UNIT IV
RANDOM VARIABLES**

1	Two dice are thrown. Let X assign to each point (a, b) in S the maximum of its numbers i.e, $X(a, b) = \max(a, b)$. Find the probability distribution. X is a random variable with $X(s) = \{1, 2, 3, 4, 5, 6\}$. Also find the mean and variance of the distribution.	[L1][CO4]	[12M]																		
2	A random variable X has the following probability function <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>X</td> <td>0</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> <td>6</td> <td>7</td> </tr> <tr> <td>P(x)</td> <td>0</td> <td>K</td> <td>2K</td> <td>2K</td> <td>3K</td> <td>K^2</td> <td>$2K^2$</td> <td>$7K^2 + K$</td> </tr> </table> <p>Determine (i) K (ii) Mean (iii) variance. (iv) if $P(X \leq K) > 1/2$, find the Minimum value of K</p>	X	0	1	2	3	4	5	6	7	P(x)	0	K	2K	2K	3K	K^2	$2K^2$	$7K^2 + K$	[L5][CO4]	[12M]
X	0	1	2	3	4	5	6	7													
P(x)	0	K	2K	2K	3K	K^2	$2K^2$	$7K^2 + K$													
3	a) Find the mean and variance of the uniform probability distribution given by $f(x) = \frac{1}{n}$ for $x = 1, 2, \dots, n$. b) If a random variable has a Probability density $f(x)$ as $f(x) = \begin{cases} 2e^{-2x}, & \text{for } x > 0 \\ 0, & \text{for } x \leq 0 \end{cases}$ Find the Probabilities that it will take on a value (i) Between 1 & 3 (ii) Greater than 0.5	[L1][CO4]	[06M]																		
4	Probability density function of a random variable X is $f(x) = \begin{cases} \frac{1}{2} \sin x, & \text{for } 0 \leq x \leq \pi \\ 0, & \text{elsewhere} \end{cases}$. Find the mean, mode and median of the distribution and also find the probability between 0 and $\frac{\pi}{2}$.	[L6][CO4]	[12M]																		
5	a) Probability density function $f(x) = \begin{cases} k(3x^2 - 1), & \text{in } -1 \leq x \leq 2 \\ 0, & \text{elsewhere} \end{cases}$. i) Find the value of k. ii) Find the probability ($-1 \leq x \leq 0$) b) The probability density function of a random variable x is $f(x) = \begin{cases} kx(x - 1); & 1 \leq x \leq 4 \\ 0; & \text{elsewhere} \end{cases}$ And $P(1 \leq x \leq 3) = \frac{28}{3}$ Find the value of k	[L1][CO4]	[06M]																		
6	For the continuous probability function $f(x) = \begin{cases} kx^2 e^{-x} & \text{when } x \geq 0 \\ 0; & \text{elsewhere} \end{cases}$ Find i) k ii) Mean iii) Variance.	[L1][CO4]	[12M]																		
7	a) Define Probability density function. b) A continuous random variable x has the distribution function $F(x) = \begin{cases} 0 & \text{if } x \leq 1 \\ k(x - 1)^4; & 1 < x \leq 3 \\ 0; & x > 3 \end{cases}$ Find the value of k and the probability density function of x.	[L1][CO4]	[02M]																		
8	a) Define Probability Distribution functions.. b) A random variable x has the following probability distribution <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>x</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> <td>6</td> </tr> <tr> <td>P(x)</td> <td>k</td> <td>3k</td> <td>5k</td> <td>7k</td> <td>9k</td> <td>11k</td> </tr> </table> Find i) k ii) Mean iii) Variance.	x	1	2	3	4	5	6	P(x)	k	3k	5k	7k	9k	11k	[L1][CO4] [L6][CO4]	[02M] [10M]				
x	1	2	3	4	5	6															
P(x)	k	3k	5k	7k	9k	11k															
9	A random variable x has the following probability distribution function <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>x</td> <td>-3</td> <td>-2</td> <td>-1</td> <td>0</td> <td>1</td> <td>2</td> <td>3</td> </tr> <tr> <td>P(x)</td> <td>k</td> <td>0.1</td> <td>k</td> <td>0.2</td> <td>2k</td> <td>0.4</td> <td>2k</td> </tr> </table> Find i) k ii) Mean iii) Variance.	x	-3	-2	-1	0	1	2	3	P(x)	k	0.1	k	0.2	2k	0.4	2k	[L6][CO4]	[12M]		
x	-3	-2	-1	0	1	2	3														
P(x)	k	0.1	k	0.2	2k	0.4	2k														
10	A random variable x has the following probability distribution function <table border="1" style="margin-left: auto; margin-right: auto;"> <tr> <td>x</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> <td>6</td> <td>7</td> <td>8</td> </tr> <tr> <td>P(x)</td> <td>k</td> <td>2k</td> <td>3k</td> <td>4k</td> <td>5k</td> <td>6k</td> <td>7k</td> <td>8k</td> </tr> </table> Find i) k ii) $P(X \leq 2)$ iii) $P(2 \leq x \leq 5)$.	x	1	2	3	4	5	6	7	8	P(x)	k	2k	3k	4k	5k	6k	7k	8k	[L1][CO4]	[12M]
x	1	2	3	4	5	6	7	8													
P(x)	k	2k	3k	4k	5k	6k	7k	8k													

UNIT V
PROBABILITY DISTRIBUTIONS AND CORRELATION

1	a) Derive mean and variance of Binomial distribution. b) 20% of items produced from a factory are defective. Find the probability that in a sample of 5 chosen at random (i) one is defective (ii) $p(1 < x < 4)$	[L3][CO5] [L6][CO5]	[06M] [06M]																																	
2	Fit a Binomial distribution to the following frequency distribution: <table border="1" style="margin-left: 40px;"> <tr> <td>x</td> <td>0</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> </tr> <tr> <td>f</td> <td>2</td> <td>14</td> <td>20</td> <td>34</td> <td>22</td> <td>8</td> </tr> </table>	x	0	1	2	3	4	5	f	2	14	20	34	22	8	[L5][CO5]	[12M]																			
x	0	1	2	3	4	5																														
f	2	14	20	34	22	8																														
3	Out of 800 families with 5 children each, how many would you expect to have (i) 3 boys (ii) 5 girls (iii) either 2 or 3 boys iv) At least one boy	[L5][CO5]	[12M]																																	
4	a) If 2% of light bulbs are defective. Find the probability that (i) At least one is defective (ii) $p(1 < x < 8)$ in a sample of 100 b) If for a poisson variate $2 P(X=0) = P(X=2)$ Find the probability that i) $P(X \leq 3)$ ii) $P(2 < X \leq 5)$ iii) $P(X \geq 3)$.	[L1][CO5] [L1][CO5]	[06M] [06M]																																	
5	Fit a Poisson distribution to the following data <table border="1" style="margin-left: 40px;"> <tr> <td>x</td> <td>0</td> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> <td>Total</td> </tr> <tr> <td>f</td> <td>142</td> <td>156</td> <td>69</td> <td>27</td> <td>5</td> <td>1</td> <td>400</td> </tr> </table>	x	0	1	2	3	4	5	Total	f	142	156	69	27	5	1	400	[L5][CO5]	[12M]																	
x	0	1	2	3	4	5	Total																													
f	142	156	69	27	5	1	400																													
6	In a sample of 1000 cases, the mean of certain test is 14 and standard deviation is 2.5. Assuming the distribution to be normal find (i) How many students score between 12 and 15. (ii) How many students score above 18? (iii) How many students score below 18?	[L6][CO5]	[12M]																																	
7	a) The probability of poisson variate taking the values 1 & 2 are equal. Find i) μ ii) $P(X \geq 1)$ iii) $P(1 < X < 4)$. b) If X is a normal variate with mean 30 and standard deviation 5. Find the probability that i) $26 \leq X \leq 40$ ii) $X \geq 45$.	[L1][CO5] [L1][CO5]	[06M] [06M]																																	
8	Calculate Correlation coefficient to the following data <table border="1" style="margin-left: 40px;"> <tr> <td>X</td> <td>10</td> <td>15</td> <td>12</td> <td>17</td> <td>13</td> <td>16</td> <td>24</td> <td>14</td> <td>22</td> <td>20</td> </tr> <tr> <td>Y</td> <td>30</td> <td>42</td> <td>45</td> <td>46</td> <td>33</td> <td>34</td> <td>40</td> <td>35</td> <td>39</td> <td>38</td> </tr> </table>	X	10	15	12	17	13	16	24	14	22	20	Y	30	42	45	46	33	34	40	35	39	38	[L5][CO5]	[12M]											
X	10	15	12	17	13	16	24	14	22	20																										
Y	30	42	45	46	33	34	40	35	39	38																										
9	Ten competitors in a musical test were ranked by the three judges A, B and C in the following order <table border="1" style="margin-left: 40px;"> <tr> <td>Ranks by A</td> <td>1</td> <td>6</td> <td>5</td> <td>10</td> <td>3</td> <td>2</td> <td>4</td> <td>9</td> <td>7</td> <td>8</td> </tr> <tr> <td>Ranks by B</td> <td>3</td> <td>5</td> <td>8</td> <td>4</td> <td>7</td> <td>10</td> <td>2</td> <td>1</td> <td>6</td> <td>9</td> </tr> <tr> <td>Ranks by C</td> <td>6</td> <td>4</td> <td>9</td> <td>8</td> <td>1</td> <td>2</td> <td>3</td> <td>10</td> <td>5</td> <td>7</td> </tr> </table> Using rank Correlation coefficient method, discuss which pair of judges has the nearest approach to common likings in music.	Ranks by A	1	6	5	10	3	2	4	9	7	8	Ranks by B	3	5	8	4	7	10	2	1	6	9	Ranks by C	6	4	9	8	1	2	3	10	5	7	[L5][CO5]	[12M]
Ranks by A	1	6	5	10	3	2	4	9	7	8																										
Ranks by B	3	5	8	4	7	10	2	1	6	9																										
Ranks by C	6	4	9	8	1	2	3	10	5	7																										
10	Find two regression equations from the following data: <table border="1" style="margin-left: 40px;"> <tr> <td>X</td> <td>10</td> <td>25</td> <td>34</td> <td>42</td> <td>37</td> <td>35</td> <td>36</td> <td>45</td> </tr> <tr> <td>Y</td> <td>56</td> <td>64</td> <td>63</td> <td>58</td> <td>73</td> <td>75</td> <td>82</td> <td>77</td> </tr> </table>	X	10	25	34	42	37	35	36	45	Y	56	64	63	58	73	75	82	77	[L1][CO5]	[12M]															
X	10	25	34	42	37	35	36	45																												
Y	56	64	63	58	73	75	82	77																												